


# *B Physics at Hadron Colliders*

Minoru TANAKA

Department of Physics

Osaka University

# 1. Introduction



- CPV in  $K \rightarrow \pi\pi$   
 $\varepsilon$  parameter
- $b \rightarrow c, u$  transition  
 $|V_{cb}|, |V_{ub}|$
- $B_d - \bar{B}_d$  mixing ( $\Delta m_{B_d}$ )  
 $|V_{td}|$
- $B_s - \bar{B}_s$  mixing ( $\Delta m_{B_s}$ )  
 $|V_{td}|$  (upper bound)
- CPA in  $B_d \rightarrow \psi K_s, \dots$   
( $A_{CP}^{mix}(b \rightarrow c\bar{c}s)$ )  
 $\sin 2\phi_1$

The unitarity triangle looks closed.

But, this is not the whole story. We want to go further.

⇒ Precision test of the standard model

⇒ New physics search

# Supersymmetry

The most attractive candidate of new physics

- Hierarchy problem: Why  $M_{\text{weak}} \ll M_{\text{GUT}}$ ?  
SUSY protects this hierarchy against radiative corrections  
For this mechanism to work,  $M_{\text{SUSY}} \sim M_{\text{weak}}$
- SUSY must be broken.  
No degenerate SUSY pair of a fermion and a boson.
- SUSY breaking must be soft.  
Cancellation of the second order divergence is required.  
Allowed soft breakings: gaugino masses, scalar masses, trilinear scalar couplings, etc.  
⇒ New sources of flavor and CP violation

B experiments will unveil these new flavor and CP violation.

# PLAN


1. Introduction
2. Hadron colliders and  $e^+e^-$  colliders
3. SUSY effects in  $b \rightarrow s$  transitions
4. Summary

## 2. Hadron colliders and $e^+e^-$ colliders

	hadron				$e^+e^-$		
	Tevatron ( $p\bar{p}$ )		LHC ( $pp$ )		KEKB PEP II	Super KEKB Super PEP II	LC
	CDF D0	BTeV	ATLAS CMS	LHCb	Belle Babar	Super Belle Super Babar	GigaZ
$\sqrt{s}$	2 TeV		14 TeV		10.6 GeV		91.2 GeV
$L$	$2 \cdot 10^{32}$		$10^{33-34}$	$2 \cdot 10^{32}$	$10^{34}$	$10^{35-36}$	$10^{34}$
$\sigma_{b\bar{b}}$	0.1 mb		0.5 mb		1.1 nb		6.6 nb
$\sigma_{\text{total}}$	75 mb		100 mb				
pileup	1.6		2-20	0.53			
	$B_s$				two missings ( $\nu\bar{\nu}$ )		$\Lambda_b$ polarization

# Physics reach: CKM

SNOWMASS 2001: Working group E2  
Electron-positron Colliders  
from the  $\rho$  to the Z



## CKM Reach

### super B factories & hadron facilities

	BTeV $10^7$ s	LHC-b $10^7$ s	BABAR BELLE (2005)	$10^{35}$ $10^7$ s	$10^{36}$ $10^7$ s	
$\sin 2\beta$	0.011	0.02	0.037	0.026	0.008	Equal
$\sin 2\alpha'$	0.05	0.05	0.14	0.1	0.032	Equal
$\gamma B_s(D_s K)$	$\sim 7^\circ$					Had
$\gamma B(DK)$	$\sim 2^\circ$		$\sim 20^\circ$		$1-2.5^\circ$	Equal
$\sin 2\chi$	0.023	0.04	--	--	--	Had
$\text{Br}(B \rightarrow \pi^0 \pi^0)$	--	--	$\sim 20\%$	14%	6%	e+e-
$V_{ub}$	--	--	$\sim 2.3\%$	$\sim 1\%$ (sys)	$\sim 1\%$ (sys)	e+e-

SBF numbers based on BABAR experience need simulations to estimate signal efficiencies & bkgds so that the comparison to BTeV/LHC-b is on an equal footing

Burdman/Butler/Shipsey/Yamamoto

(Table compiled by SBF E2 subgroup & E2 convenors)


45

$$\text{Belle } \sin 2\phi_1 = 0.719 \pm 0.074 \pm 0.035$$

$$\text{BaBar } \sin 2\beta = 0.741 \pm 0.067 \pm 0.033$$

# Physics reach: rare decays

SNOWMASS 2001: Working group E2  
Electron-positron Colliders  
from the  $\phi$  to the Z



## Rare B decay Reach super B fac. & hadron facilities

All numbers are per year except current B factories (cumulative Through 2005)

Best measurement ●

Greater reach?

Hadronic:  $B \rightarrow \mu\mu$

$e^+e^-$ :  $s\nu\nu, \tau\nu, \gamma\gamma, K^*\gamma$

Including additional rare  $\Delta b$  and  $B_s$  modes not in Table

$10^{35}$  reach < BTeV/LHCb

$10^{36}$  reach ~ BTeV/LHCb

(Table compiled by SBF E2 subgroup & E2 convenors)

Burdman/Butler/Shipsev/Yamamoto

Decay Mode (Br. Ratio)	HADRONIC EXP.			B-FACTORY		
	CDF/D0 (2 fb <sup>-1</sup> )	BTeV/LHC-b (10 <sup>7</sup> s)	ATLAS/CMS (1 Year)	BABAR/BELLE (0.5 ab <sup>-1</sup> )	10 <sup>35</sup> (1 ab <sup>-1</sup> )	10 <sup>36</sup> (10 ab <sup>-1</sup> )
$B \rightarrow X_s \gamma$ (3.29 ± .21 ± .21) × 10 <sup>-4</sup> With B Tags				11K 1.7K	22K 3.4K	220K 34K
$B \rightarrow K^* \gamma$ (3 - 8) × 10 <sup>-5</sup> $\delta(A_{CP})$	170/-	27K/24K 0.01		6K 0.02	12K 0.01	120K < 0.01
$B \rightarrow X_s \nu \bar{\nu}$ (4.1 ± 0.9) × 10 <sup>-5</sup>				8	16	160
$B \rightarrow K^* \nu \bar{\nu}$ (5 × 10 <sup>-6</sup> )				1.5	3	30
$B \rightarrow X_s \mu^+ \mu^-$ (6.0 ± 1.5) × 10 <sup>-6</sup>		7.2K/-		300	600	6K
$B \rightarrow X_s e^+ e^-$				350	700	7K
$B \rightarrow K^* \mu^+ \mu^-$ (2 ± 1) × 10 <sup>-6</sup>	61/60-150	4.4 k /4.5k	665/4.2K	120	240	2.4K
$B \rightarrow K^* e^+ e^-$				150	300	3K
$B_d^0 \rightarrow \tau^+ \tau^-$ (10 <sup>-7</sup> )				< 10 <sup>-5</sup>	< 5 × 10 <sup>-6</sup>	< 10 <sup>-6</sup>
$B^0 \rightarrow \mu^+ \mu^-$ $B_s$ (10 <sup>-9</sup> ) $B_d$ (8 × 10 <sup>-11</sup> )	5/1.5-6 0/0	10/11 2/2	9/7 0.7/20	< 10 <sup>-8</sup>	< 5 × 10 <sup>-9</sup>	< 10 <sup>-9</sup>
$B_d^0 \rightarrow e^+ e^-$ (10 <sup>-14</sup> )				< 10 <sup>-8</sup>	< 5 × 10 <sup>-9</sup>	< 10 <sup>-9</sup>
$B \rightarrow \tau \nu$ (5 × 10 <sup>-5</sup> )				17	34	350
$B \rightarrow \mu \nu$ (1.6 × 10 <sup>-7</sup> )				8	16	150
$B^0 \rightarrow \gamma \gamma$ (10 <sup>-8</sup> )				0.4	0.8	8

### 3. SUSY effects in $b \rightarrow s$ transitions

The large 2–3 mixing in the  $\nu$  oscillation

- $B \rightarrow X_s \gamma, B \rightarrow K^* \gamma$
  - $B \rightarrow X_s l^+ l^-, B \rightarrow K^{(*)} l^+ l^-$
  - $B \rightarrow X_s \nu \bar{\nu}, B \rightarrow K^{(*)} \nu \bar{\nu}$
  - $B \rightarrow \phi K_S$
  - $B \rightarrow \eta' K_S$
  - $B \rightarrow K \pi$
  - $B_s$  mixing ( $\Delta m_{B_s}, \Delta \Gamma_{B_s}$ )
  - $B_s \rightarrow \mu^+ \mu^-$
- etc.

# Soft *SUSY* breakings

Scalar masses, scalar trilinear couplings

⇒ New sources of flavor and CP violation

- Squark mass matrix ( $6 \times 6$ )

$$\tilde{M}_q^2 = \begin{pmatrix} \tilde{M}_{q,LL}^2 & \tilde{M}_{q,LR}^2 \\ \tilde{M}_{q,LR}^{2\dagger} & \tilde{M}_{q,RR}^2 \end{pmatrix}, \quad (q = u, d)$$

- Quark mass matrix ( $3 \times 3$ )

$$V_L^q M_q V_R^{q\dagger} = \hat{M}_q \text{ (diagonal)}$$

FV is caused by off-diagonal

$$(\delta_M^{qN})_{ij} \equiv \frac{(V_M^q \tilde{M}_{q,MN}^2 V_N^{q\dagger})_{ij}}{\tilde{m}^2}$$

- $K-\bar{K}$  mixing

$$\sqrt{(\delta_{LL}^d)_{12}(\delta_{RR}^d)_{12}} < 0.006, \quad (\delta_{MM}^d)_{12} < 0.05$$

$$\sqrt{(\delta_{LL}^u)_{12}(\delta_{RR}^u)_{12}} < 0.04, \quad (\delta_{MM}^u)_{12} < 0.1$$

- $B-\bar{B}$  mixing

$$\sqrt{(\delta_{LL}^d)_{13}(\delta_{RR}^d)_{13}} < 0.04, \quad (\delta_{MM}^d)_{13} < 0.1$$

- $b \rightarrow s\gamma$

$$(\delta_{LR}^d)_{23} < 0.03 \quad (\tilde{m} \sim 400\text{GeV}) \quad \text{Kane et al., hep-ph/0212092}$$

$$\Rightarrow (\delta_{MN}^q)_{ij} < 0.01 \sim 0.1 \quad (\tilde{m} \sim 1\text{TeV})$$

We need some mechanism to suppress  $\delta$ 's.

Note: no constraint on  $(\delta_{LL,RR}^d)_{23}$

# Possible mechanisms

Several ideas in the market

- Universal soft breakings  
Gravity mediation of SUSY breaking (SUGRA),  
Gauge mediation  
Phenomenology depends on the scale of SUSY breaking and extra interaction below it (GUT,  $\nu$  Yukawa).
- Flavor symmetry  
Must explain the fermion masses and mixings.  
Depends on the breaking sector.
- Alignment
- Effective SUSY

SUSY breaking mechanism  $\iff \delta's \iff$  B physics

# $B_s - \bar{B}_s$ mixing: $|V_{td}|$

$\Delta m_{B_s} = 2|M_{12}(B_s)|$ : box diagram

- Hadronic uncertainty

$$f_{B_s} \sqrt{B_{B_s}} = 276 \pm 38 \text{ MeV (lattice)}$$

Taking a ratio  $\Delta m_{B_s} / \Delta m_{B_d}$  reduces the hadronic uncertainty.  $\Leftarrow$  SU(3) symmetry.

$$f_{B_s} \sqrt{B_{B_s}} / f_{B_d} \sqrt{B_{B_d}} = 1.18 \pm 0.04_{-0}^{+0.12} \text{ (lattice)}$$

But, the SU(3) breaking may be larger.

# $B_s-\bar{B}_s$ mixing: $|V_{td}|$

$\Delta m_{B_s} = 2|M_{12}(B_s)|$ : box diagram


- Hadronic uncertainty

$$f_{B_s} \sqrt{B_{B_s}} = 276 \pm 38 \text{ MeV}$$

Taking a ratio  $\Delta m_{B_s}/\Delta m_{B_d}$   
uncertainty.  $\Leftarrow$  SU(3) sym

$$f_{B_s} \sqrt{B_{B_s}} / f_{B_d} \sqrt{B_{B_d}} =$$

But, the SU(3) breaking n



# $B_s-\bar{B}_s$ mixing: $|V_{td}|$

$\Delta m_{B_s} = 2|M_{12}(B_s)|$ : box diagram

- Hadronic uncertainty

$$f_{B_s} \sqrt{B_{B_s}} = 276 \pm 38 \text{ MeV (lattice)}$$

Taking a ratio  $\Delta m_{B_s} / \Delta m_{B_d}$  reduces the hadronic uncertainty.  $\Leftarrow$  SU(3) symmetry.

$$f_{B_s} \sqrt{B_{B_s}} / f_{B_d} \sqrt{B_{B_d}} = 1.18 \pm 0.04_{-0}^{+0.12} \text{ (lattice)}$$

But, the SU(3) breaking may be larger.

- Possible new physics contribution

$$M_{12}(B_s) = M_{12}(B_s)|_{\text{SM}} + M_{12}(B_s)|_{\text{NP}}$$

The effect might be large.

cf. large 2–3 mixing in the  $\nu$  sector

# $\Delta m_{B_s}$

- Experiment

$$\Delta m_{B_s} > 14.9 \text{ ps}^{-1} \text{ (95\% CL)}$$

$$\text{cf. } \Delta m_{B_d} = 0.489 \pm 0.008 \text{ ps}^{-1}$$

- SM prediction

$$\Delta m_{B_s} = 17.3_{-0.7}^{+1.5} \text{ ps}^{-1}$$

- SUSY contribution

Could be large.

$$(\delta_{LL}^q) \sim 0.5 \Rightarrow \Delta m_{B_s} \sim 30 \text{ ps}^{-1}$$


- Experimental reach

$$\text{CDF } 38 \sim 48 \text{ ps}^{-1}, \text{ BTeV } \sim 49 \text{ ps}^{-1}$$

$$\text{ATLAS } 30 \text{ ps}^{-1}, \text{ CMS } 26 \text{ ps}^{-1}, \text{ LHCb } 48 \text{ ps}^{-1}$$

# SUSY effect in $\Delta m_{B_s}$

mSUGRA  $\sim O(0.01)$ , SU(5)  $\sim O(0.01)$ , U(2)  $\sim O(1)$




T. Goto, Y. Okada, Y. Shimizu, T. Shindou, M.T.

Phys. Rev. D 66, 035009 (2002)

# Correlations


$\Delta m_{B_s} / \Delta m_{B_d}$ ,  $A_{CP}^{\text{mix}}(B \rightarrow \psi K_S)$ , and  $\phi_3$



T. Goto, Y. Okada, Y. Shimizu, T. Shindou, M.T.

Phys. Rev. D 66, 035009 (2002)

# SUSY effect in $\Delta m_{B_s}$ (cont'd)



$\Delta m_{B_s}$

$B \rightarrow \phi K_S$  implied.

Ciuchini et al.  
hep-ph/0212397

# $B_s \rightarrow l^+l^-$ ( $b \rightarrow sl^+l^-$ )

- Experiment

$$\text{BR}(B_s \rightarrow \mu^+\mu^-) < 2.6 \times 10^{-6} \text{ (CDF, 95\% CL)}$$

- SM prediction

$$\text{BR}(B_s \rightarrow \mu^+\mu^-) = (3.1 \pm 1.4) \times 10^{-9}$$

- Experimental reach

$$\text{CDF} \sim 1 \times 10^{-8}$$

ATLAS, CMS (3yrs)  $\sim$  SM

LHCb (1yr)  $\sim$  SM

# $B_s \rightarrow l^+ l^-$ (cont'd)

- SUSY contribution  
FCNC via neutral Higgs ( $h$ ,  $H$  and  $A$ ) exchange


Huang and Yan, PLB442 209(1998); Huang, Liao and Yan, PRD59 011701(1998); Yan et al., PRD62 094023(2000); Huang et al., PRD63 114021(2001), PRD64 059902(E) (2001); Choudhury and Gaur, PLB451 86 (1999); Babu and Kolda, PRL84(2000); Chankowski and Sławaianowska, PRD63 054012 (2001); Dedes, Dreiner and Nierste, PRL87 251804 (2001); Dedes and Pilaftsis, PRD67 015012 (2003); Bobeth et al., PRD64 074014 (2001), PRD66 074021 (2002); Mizukoshi, Tata and Wang, PRD66 115003 (2002).

$$A(B_s \rightarrow \mu^+ \mu^-) \sim \tan^3 \beta \quad \text{as } \tan \beta (= \frac{v_u}{v_d}) \rightarrow \infty$$

Does not decouple as  $M_{\text{SUSY}} \rightarrow \infty$ .

Decouples as  $m_A \rightarrow \infty$ .

# $B_S \rightarrow l^+l^-$ (cont'd)



MFV

$$m_{\tilde{t}_1} = 120 \text{ GeV}$$

$$\theta_{\tilde{t}} \approx -45^\circ$$

Bobeth et al.  
PRD64 074014  
(2001)

## 4. Summary

- $B_s$  physics is one of the advantages of hadron B experiments.  
 $B_s$  mixing measurement by CDF is important.
- Modes with two (or more) missing and modes without charged tracks are difficult for hadron B experiments.
- A dedicated hadron B experiment  $\sim$  A  $10^{36}$  B factory
- SUSY effects could be found in the  $b \rightarrow s$  transitions.
- There are lots of other interesting modes,  
 $B_s \rightarrow J/\psi\phi$ ,  $B_d \rightarrow \rho\pi$ ,  $B \rightarrow DD$ ,  $B \rightarrow D^{(*)}\pi$ , etc.